

Science behind the danger of blue light to the retina

By: Shelby Temple, BSc, MSc, PhD

Light is part of the electromagnetic radiation spectrum and while vital for life on earth and necessary for vision it carries energy that can be damaging, specifically to the region of the retina called the macula where light is best focussed.

There is a wealth of evidence^{1,2,3,4} that it is the blue end of the visible spectrum of light that is most dangerous to the retina. Ultraviolet and infrared are almost entirely filtered out by the lens, cornea and vitreous⁵ and therefore do not reach the retina. Short-wavelength, high energy visible, blue light (380-500 nm) has enough energy per photon (>2.5 eV) to cause damage to cells and tissues known as photochemical damage^{2,3,6}. Photochemical damage typically manifests through the creation of reactive oxygen species and free radicals, which can damage proteins, cell membranes, RNA, DNA and mitochondrial DNA^{7,8,9}. The accumulation of damage to DNA and mitochondrial DNA has been linked to the process of aging (free radical theory of aging)¹⁰ and is thought to play a role in the aetiology of age-related macular degeneration¹¹.

Research, showing evidence of photochemical damage, has been performed at various levels from molecular^{12,13,14}, cellular^{15,16,17,18,19,20} and tissue^{21,22,23} all the way to whole animals^{24,25,26,27,28,29,30} including primates^{31,32,33} and humans^{34,35,36,37,38}. These studies consistently provide evidence to support the argument that short-term high-intensity blue light causes photochemical damage, and that blue light is more dangerous than longer wavelengths. Evidence for long-term low intensity (type I) photochemical damage leading to age related macular degeneration (AMD) in humans has been obtained from epidemiological studies^{3,39,40,41,42,43,44,45,46,47}. And, while there are some epidemiological studies that have failed to detect a correlation between long-term light exposure and AMD^{48,49}, it is probable that this is due to the multifactorial nature of aging and large number of uncontrolled variables involved in such studies.

Controlled studies of long-term photochemical damage are not possible in humans due to the length of time required for such studies (>50 years), however, research with monkeys, which have eyes that are structurally identical to our own, has shown that blue light can cause early signs of macular degeneration (lipofuscin and drusen accumulation)^{33,50,51,52}. In these controlled studies, the maculae were exposed to greater amounts of blue light through a reduction in macular pigments (lutein, zeaxanthin and meso-zeaxanthin), removed by eliminating carotenoids from the monkeys' diet, compared to control groups that had a normal diet. These studies have shown that in the absence of blue light protection early signs of AMD can occur at half the age (human age equivalent of 35 instead of 65 years of age)⁵². Humans also have macular pigments that are our natural blue light defence system.

The French National Health and Safety Authority (ANSES) released an extensive report in April of 2019 addressing the issue of blue light, specifically in relation to LED lighting and risks of long-term photochemical damage and AMD. In that report, they recognise that the

standards set on the intensity of blue light emitted from LED lights is based on Type II (short term) retinal damage and is too high given the potential long-term exposure that many people are experiencing with these commercially available light sources.

The International Organisation for Standardisation (ISO) published a technical report (ISO-TR-20772), in late 2018, as guidance for establishing unified standards against which blue light filtering lenses can be compared. This was done in recognition of the benefits they offer in relation to long-term photochemical damage to the macula by blue light. The report identified 380-455 nm as the wavelengths of greatest concern³⁶. It represents the first step in a process that was previously undertaken for establishing the UV400 certification for sunglasses. In the ISO report, they identify the strong risk factors for accelerated aging of the macula as age, smoking, low macular pigments and genetics.

The ISO report also recognises the value of allowing the “good” blue light 455 - 500 nm to reach the eye. This is because a part of the blue light spectrum 455 - 500 nm helps entrain our circadian rhythms (sleep/wake cycle) through a pigment called melanopsin, which has a peak photosensitivity between 460-480 nm^{4,53,54,55}. This is relevant to front of eye methods of filtering blue light reaching the eye (interocular lenses, contact lenses and glasses) because they intervene before light reaches the retina, whereas macular pigments are located in the retina itself and therefore only filter blue light in the macula thus allowing blue light to reach the rest of the retina where melanopsin can be activated.

There is a great deal of misinformation about the dangers of blue light and many sources confuse blue light with light exclusively from electronic devices. As such governing bodies in optometry in the UK (e.g. GOC and AOP) are making statements that are misleading e.g. “There is no evidence that visible blue light causes eye disease in humans”, which is simply not true (as above). Knowledge about the danger of blue light is hindered by claims that there is a lack of “strong” or “high-quality” evidence to support the contention that blue light causes retinal damage, because people uneducated in the field think that this means that there is “no evidence” that blue light is dangerous (but see above).

It is important to recognise that the lack of evidence does not mean that the statement is untrue, but rather that the type of evidence that these organizations are looking for (e.g. double-blind placebo-controlled studies in humans) is not yet, and may never be, available. The same argument was made by the tobacco industry for many years until the combination of different types of evidence (exactly like we have now for blue light) were brought together in the US Surgeon General’s report in 1964⁵⁶. The question then is: did smoking cause cancer before 1964? Of course, it did. Sadly, if governing bodies had responded more quickly to the evidence that had been accumulating for over 30 years, millions of lives could have been saved. We should not risk millions of people losing their vision prematurely because we ignored the evidence we have already.

For blue light, the type of study expected by some of the governing bodies will never exist because of the nature of age-related macular degeneration (AMD), which is caused by a lifelong accumulation of oxidative photochemical damage due to free radicals (reactive oxygen species). As such a controlled study in humans to prove the efficacy of blue light filtering lenses would require greater than 50 years and control measures that would not be

acceptable for a human study. However, there is a great deal of evidence to support the link between blue light and AMD (see citations).

In the absence of a long-term controlled study in humans showing a causative link between blue light and retinal disease, it would be advisable, based on the evidence that does exist, for eye care practitioners to recommend that patients take precautionary action by limiting their exposure (both time and intensity) to short-wavelength blue light. This advice can include limiting time in direct sunlight*, wearing hats*, sunglasses*, ophthalmic lenses that reduce the proportion of blue light, as well as improving diet by consuming more dark and brightly coloured fruits and vegetables* or supplements that contain macular pigments.

*These are actions recommended by the Macular Society to help prevent long-term retinal damage.

I would urge anyone interested in this topic to learn more by reading reviews by Boulton et al. 2001, Tosini et al. 2016 and Young 1988, and looking at Figure 2 in Fletcher et al. 2008 who showed that the odds ratio of getting AMD for people with low macular pigments (our natural protection against blue light) combined low vitamin C and E was nearly as high (2.61-3.72) as the odds ratio for smokers (4.0). This is relevant because only 17% of the UK population smoke but 25-30% of UK population have low macular pigment levels. There are many other good papers below, and this is just the tip of the iceberg with respect to research in this area.

Citations

- 1 Glickman, R. D. Phototoxicity to the retina: Mechanisms of damage. *Int. J. Toxicol.* **21**, 473-490, doi:10.1080/10915810290169909 (2002).
- 2 Boulton, M., Rozanowska, M. & Rozanowski, B. Retinal photodamage. *J. Photochem. Photobiol. B Biol.* **64**, 144-161, doi:10.1016/s1011-1344(01)00227-5 (2001).
- 3 Young, R. W. Solar radiation and age-related macular degeneration. *Surv. Ophthalmol.* **32**, 252-269, doi:10.1016/0039-6257(88)90174-9 (1988).
- 4 ISO, I. O. f. S. Vol. TR 20772:2018 34 (ISO copyright office, Geneva, 2018).
- 5 Schwartz, S. G., Gombos, D. S. & Schneider, S. in *Duane's Ophthalmology* eds W. Tasman & E.A. Jaeger (Lippincott Williams & Wilkins, 2011).
- 6 Wu, J. M., Seregard, S. & Algvere, P. V. Photochemical damage of the retina. *Surv. Ophthalmol.* **51**, 461-481, doi:10.1016/j.survophthal.2006.06.009 (2006).
- 7 Godley, B. F., Shamsi, F. A., Liang, F. Q., Jarrett, S. G., Davies, S. & Boulton, M. Blue light induces mitochondrial DNA damage and free radical production in epithelial cells. *J. Biol. Chem.* **280**, 21061-21066, doi:10.1074/jbc.M502194200 (2005).
- 8 Lockwood, D. B., Wataha, J. C., Lewis, J. B., Tseng, W. Y., Messer, R. L. W. & Hsu, S. D. Blue light generates reactive oxygen species (ROS) differentially in tumor vs. normal epithelial cells. *Dent. Mater.* **21**, 683-688, doi:10.1016/j.dental.2004.07.022 (2005).
- 9 Pflaum, M., Kielbassa, C., Garmyn, M. & Epe, B. Oxidative DNA damage induced by visible light in mammalian cells: extent, inhibition by antioxidants and genotoxic effects. *Mutation Research-DNA Repair* **408**, 137-146, doi:10.1016/s0921-8777(98)00029-9 (1998).
- 10 Grimm, A. & Eckert, A. Brain aging and neurodegeneration: from a mitochondrial point of view. *J. Neurochem.* **143**, 418-431, doi:10.1111/jnc.14037 (2017).
- 11 Jarrett, S. G., Lewin, A. S. & Boulton, M. E. in *Studies on retinal and choroidal disorders Oxidative Stress in Applied Basic Research and Clinical Practice* (ed R.D. Stratton et al.) Ch. 10, 203-239 (2012).
- 12 Nakanishi-Ueda, T. et al. Blue LED light exposure develops intracellular reactive oxygen species, lipid peroxidation, and subsequent cellular injuries in cultured bovine retinal pigment epithelial cells. *Free Radic. Res.* **47**, 774-780, doi:10.3109/10715762.2013.829570 (2013).
- 13 Ratnayake, K., Payton, J. L., Lakmal, O. H. & Karunaratne, A. Blue light excited retinal intercepts cellular signaling. *Sci. Rep.-UK* **8**, 16, doi:10.1038/s41598-018-28254-8 (2018).
- 14 Alaimo, A. et al. Toxicity of blue led light and A2E is associated to mitochondrial dynamics impairment in ARPE-19 cells: implications for age-related macular degeneration. *Arch. Toxicol.* **93**, 1401-1415, doi:10.1007/s00204-019-02409-6 (2019).
- 15 Sparrow, J. R., Nakanishi, K. & Parish, C. A. The lipofuscin fluorophore A2E mediates blue light-induced damage to retinal pigmented epithelial cells. *Invest. Ophthalmol. Vis. Sci.* **41**, 1981-1989 (2000).
- 16 Roehlecke, C., Schaller, A., Knels, L. & Funk, R. H. W. The influence of sublethal blue light exposure on human RPE cells. *Mol. Vis.* **15**, 1929-1938 (2009).
- 17 Knels, L. et al. Blue light stress in retinal neuronal (R28) cells is dependent on wavelength range and irradiance. *Eur. J. Neurosci.* **34**, 548-558, doi:10.1111/j.1460-9568.2011.07790.x (2011).
- 18 Yoshida, A. et al. Reactive oxygen species production in mitochondria of human gingival fibroblast induced by blue light irradiation. *Journal of Photochemistry and Photobiology B-Biology* **129**, 1-5, doi:10.1016/j.jphotobiol.2013.09.003 (2013).
- 19 Sparrow, J. R., Miller, A. S. & Zhou, J. L. Blue light-absorbing intraocular lens and retinal pigment epithelium protection in vitro. *J. Cataract Refract. Surg.* **30**, 873-878, doi:10.1016/j.jcrs.2004.01.031 (2004).
- 20 Davies, S. et al. Photocytotoxicity of lipofuscin in human retinal pigment epithelial cells. *Free Radic. Biol. Med.* **31**, 256-265, doi:10.1016/s0891-5849(01)00582-2 (2001).

- 21 Youn, H. Y., Chou, B. R., Cullen, A. P. & Sivak, J. G. Effects of 400 nm, 420 nm, and 435.8 nm radiations on cultured human retinal pigment epithelial cells. *Journal of Photochemistry and Photobiology B-Biology* **95**, 64-70, doi:10.1016/j.jphotobiol.2009.01.001 (2009).
- 22 Wihlmark, U., Wrigstad, A., Roberg, K., Nilsson, S. E. G. & Brunk, U. T. Lipofuscin accumulation in cultured retinal pigment epithelial cells causes enhanced sensitivity to blue light irradiation. *Free Radic. Biol. Med.* **22**, 1229-1234, doi:10.1016/s0891-5849(96)00555-2 (1997).
- 23 Mignon, C., Uzunbajakava, N. E., Castellano-Pellicena, I., Botchkareva, N. V. & Tobin, D. J. Differential response of human dermal fibroblast subpopulations to visible and near-infrared light: Potential of photobiomodulation for addressing cutaneous conditions. *Lasers Surg. Med.* **50**, 859-882, doi:10.1002/lsm.22823 (2018).
- 24 Nash, T. R. *et al.* Daily blue-light exposure shortens lifespan and causes brain neurodegeneration in *Drosophila*. *Npj Aging and Mechanisms of Disease* **5**, 8, doi:10.1038/s41514-019-0038-6 (2019).
- 25 Vicente-Tejedor, J. *et al.* Removal of the blue component of light significantly decreases retinal damage after high intensity exposure. *PLoS ONE* **13**, 18, doi:10.1371/journal.pone.0194218 (2018).
- 26 Krigel, A. *et al.* Light-induced retinal damage using different light sources, protocols and rat strains reveals LED phototoxicity. *Neuroscience* **339**, 296-307, doi:10.1016/j.neuroscience.2016.10.016 (2016).
- 27 Sisson, T. R. C., Glauser, S. C., Glauser, E. M., Tasman, W. & Kuwabara, T. Retinal changes produced by phototherapy. *J. Pediatr.* **77**, 221-+, doi:10.1016/s0022-3476(70)80327-4 (1970).
- 28 Marshall, J., Mellerio, J. & Palmer, D. A. Damage to pigeon retinae by moderate illumination from fluorescent lamps. *Exp. Eye Res.* **14**, 164-8, doi:10.1016/0014-4835(72)90063-2 (1972).
- 29 Specht, S., Leffak, M., Darrow, R. M. & Organisciak, D. T. Damage to rat retinal DNA induced in vivo by visible light. *Photochem. Photobiol.* **69**, 91-98, doi:10.1562/0031-8655(1999)069<0091:dtrrdi>2.3.co;2 (1999).
- 30 Noell, W. K., Walker, V. S., Kang, B. S. & Berman, S. Retinal damage by light in rats. *Invest. Ophthalmol. Vis. Sci.* **5**, 450-473 (1966).
- 31 Ham, W. T., Ruffolo, J. J., Mueller, H. A., Clarke, A. M. & Moon, M. E. Histologic analysis of photochemical lesions produced in rhesus retina by short-wavelength light. *Invest. Ophthalmol. Vis. Sci.* **17**, 1029-1035 (1978).
- 32 Ham, W. T., Mueller, H. A. & Sliney, D. H. Retinal sensitivity to damage from short wavelength light. *Nature* **260**, 153-155, doi:10.1038/260153a0 (1976).
- 33 Barker, F. M. *et al.* Nutritional manipulation of primate retinas, V: effects of lutein, zeaxanthin, and n-3 fatty acids on retinal sensitivity to blue-light-induced damage. *Invest. Ophthalmol. Vis. Sci.* **52**, 3934-3942, doi:10.1167/iovs.10-5898 (2011).
- 34 Tso, M. O. & La Piana, F. G. The human fovea after sungazing. *Trans Sect Ophthalmol Am Acad Ophthalmol Otolaryngol.* **79**, OP788-795 (1975).
- 35 Solley, W. A. & Sternberg, P. Retinal phototoxicity. *Int. Ophthalmol. Clin.* **39**, 1-12, doi:10.1097/00004397-199903920-00002 (1999).
- 36 Kleinmann, G., Hoffman, P., Schechtman, E. & Pollack, A. Microscope-induced retinal phototoxicity in cataract surgery of short duration. *Ophthalmology* **109**, 334-338, doi:10.1016/s0161-6420(01)00924-1 (2002).
- 37 Niwano, Y., Iwasawa, A., Tsubota, K., Ayaki, M. & Negishi, K. Protective effects of blue light-blocking shades on phototoxicity in human ocular surface cells. *BMJ Open Ophthalmol.* **4**, 6, doi:10.1136/bmjophth-2018-000217 (2019).
- 38 Organisciak, D. T. & Vaughan, D. K. Retinal light damage: Mechanisms and protection. *Prog. Retin. Eye Res.* **29**, 113-134, doi:10.1016/j.preteyeres.2009.11.004 (2010).
- 39 Taylor, H. R., Muñoz, B., West, S., Bressler, N. M., Bressler, S. B. & Rosenthal, F. S. Visible light and risk of age-related macular degeneration. *Trans. Am. Ophthalmol. Soc.* **88**, 163-173 (1990).
- 40 Sui, G. Y. *et al.* Is sunlight exposure a risk factor for age-related macular degeneration? A systematic review and meta-analysis. *Br. J. Ophthalmol.* **97**, 389-394, doi:10.1136/bjophthalmol-2012-302281 (2013).
- 41 Cruickshanks, K. J., Klein, R. & Klein, B. E. K. Sunlight and age-related macular degeneration - The Beaver Dam eye study. *Arch. Ophthalmol.* **111**, 514-518, doi:10.1001/archophth.1993.01090040106042 (1993).
- 42 Klein, R., Klein, B. E. K., Jensen, S. C. & Cruickshanks, K. J. The relationship of ocular factors to the incidence and progression of age-related maculopathy. *Arch. Ophthalmol.* **116**, 506-513, doi:10.1001/archophth.116.4.506 (1998).
- 43 Fletcher, A. E. *et al.* Sunlight exposure, antioxidants, and age-related macular degeneration. *Arch. Ophthalmol.* **126**, 1396-1403, doi:10.1001/archophth.126.10.1396 (2008).
- 44 Prevention of health Damage by the Sun - Relationship Prevention in the City and in the Countryside: the Basis Paper of the UV Protection Alliance. *Bundesgesundheitsblatt-Gesundheitsforschung-Gesundheitsschutz* **60**, 1153-1160, doi:10.1007/s00103-017-2619-5 (2017).
- 45 Vojnikovic, B., Njiric, S., Coklo, M. & Spanjol, J. Ultraviolet sun radiation and incidence of age-related macular degeneration on Croatian island Rab. *Coll. Antropol.* **31**, 43-44 (2007).
- 46 Vojnikovic, B. & Vojnikovic, D. Benedictine Monastery on the Island of Rab Hides the Mystery of Sunlight Influence on Development of Macular Degeneration (AMD). Is a Harmful Sunlight, Risk Factor or Cause of Macular Degeneration? *Coll. Antropol.* **34**, 5-8 (2010).
- 47 Modenese, A. & Gobba, F. Macular degeneration and occupational risk factors: a systematic review. *Int. Arch. Occup. Environ. Health* **92**, 1-11, doi:10.1007/s00420-018-1355-y (2019).
- 48 Bai, Z. L., Ren, B. C., Yang, J. G., He, Y. A., Chen, L. & Sun, N. X. Epidemiological investigation on age related macular degeneration in rural area of Shaanxi Province, China. *International Journal of Ophthalmology* **1**, 77-84 (2008).
- 49 Delcourt, C. *et al.* Light exposure and the risk of age-related macular degeneration - The Pathologies Oculaires Liees a l'Age (POLA) study. *Arch. Ophthalmol.* **119**, 1463-1468 (2001).
- 50 Koo, E., Neuringer, M. & SanGiovanni, J. P. Macular xanthophylls, lipoprotein-related genes, and age-related macular degeneration. *Am. J. Clin. Nutr.* **100**, 336S-346S, doi:10.3945/ajcn.113.071563 (2014).
- 51 McGill, T. J., Renner, L. M. & Neuringer, M. Elevated Fundus Autofluorescence in Monkeys Deficient in Lutein, Zeaxanthin, and Omega-3 Fatty Acids. *Invest. Ophthalmol. Vis. Sci.* **57**, 1361-1369, doi:10.1167/iovs.15-18596 (2016).
- 52 McGill, T. in *Brain and Ocular Nutrition* (ed John Nolan) (Cambridge, 2018).
- 53 Brainard, G. C. *et al.* Action spectrum for melatonin regulation in humans: Evidence for a novel circadian photoreceptor. *J. Neurosci.* **21**, 6405-6412 (2001).
- 54 Hattar, S., Liao, H. W., Takao, M., Berson, D. M. & Yau, K. W. Melanopsin-containing retinal ganglion cells: Architecture, projections, and intrinsic photosensitivity. *Science* **295**, 1065-1070, doi:DOI 10.1126/science.1069609 (2002).
- 55 Tosini, G., Ferguson, I. & Tsubota, K. Effects of blue light on the circadian system and eye physiology. *Mol. Vis.* **22**, 61-72 (2016).
- 56 US Surgeon General. Smoking and Health: Report of the advisory committee to the surgeon general of the public health service. Report No. 1103, 386 (Washington DC, 1964).